

| ANTIINFECIOSO | VIAS DE ADM. | | | PRESENTACION | DILUYENTES PARA RECONSTITUCIÓN | VOL. DIL (ml). | CONC. FINAL (mg/ml) | ESTABILIDAD | | VEHIC. de INF. | | VOL de INF. (ml) | CONCENT. INFUSIÓN (mg/ml) | TIEMPO de ADMINIST |
|-------------------------------|--------------|-------|-------|--------------|--------------------------------|----------------|---------------------|-------------|--------|----------------|-------|------------------|---------------------------|--------------------|
| | IM | IVdir | IVinf | | | | | 25 °C | 2-8 °C | SF | Dx 5% | | | |
| ACICLOVIR | | | • | FA 500 mg | AD | 10,0 | 50 | 12 hs | | • | • | 50 – 100 | 5 – 10 | ≥ 1 h. |
| AMIKACINA | • | | | FA 100 mg | | | | | | | | | | |
| | | | • | FA 500 mg | | | | | | • | • | 100 – 200 | 2,5 – 5 | 30 – 60' |
| AMPICILINA | • | | | FA 1g | AD | 3,5 | 250 | 1 h | | | | | | |
| | | • | | FA 1g | AD | 10,0 | 100 | 1 h | 4 hs | | | | | 10-15' |
| | | | • | FA 1g | AD | 10,0 | 100 | 1 h | 4 hs | • | | 50 – 100 | 10 – 20 | 15-30' |
| AMPI-SULBACTAM | • | | | FA 1,5g | AD- Lidoc. 1% | 3,2 | 250 | 1 h | | | | | | |
| | | • | | FA 1,5g | AD | 3,2 | 250 | 1 h | 4 hs | | | | | 10-15' |
| | | | • | FA 1,5g | AD | 3,2 | 250 | 1 h | 4 hs | • | | 50-100 | 10 – 20 | 15-30' |
| ANFOTERICINA | | | • | FA 50mg | AD-Dx 5% | 10,0 | 5 | 24 hs | 7 ds | | • | 500 | 0,1 | □ 6 hs |
| ANFOTERICINA LIPOSOMAL | | | • | FA 50mg | AD | 12,0 | | 24 hs | | | • | 25-100 | 2-0,2 | 4-6 hs |
| CEFAZOLINA | • | | | FA 1g | AD-SF | 2,5 | | 24 hs | 10 ds | | | | | |
| | | • | | FA 1g | AD | 5,0-10,0 | 100-200 | 24 hs | 10 ds | | | | | 3-5' |
| | | | • | FA 1g | AD | 5,0-10,0 | 100-200 | 24 hs | 10 ds | • | • | 50-100 | 10 – 20 | 30' |
| CEFALOTINA | • | | | FA 1g | AD | 4,0 | 225 | 24 hs | 4 ds | | | | | |
| | | • | | FA 1g | AD – SF-Dx 5% | 10,0 | 100-200 | 24 hs | 4 ds | | | | | 3-5' |
| | | | • | FA 1g | AD – SF-Dx 5% | 10,0 | 100-200 | 24 hs | 4 ds | • | • | 50-100 | 10 – 20 | 30' |
| CEFEPIME | • | | | FA 2 g | AD- SF-Dx 5% | 7,0 | 280 | | | | | | | |
| | | • | | FA 2 g | AD – SF-Dx 5% | 20,0 | 100 | 24 hs | 7 ds | | | | | 3-5' |
| | | | • | FA 2 g | AD | 10,0 | 200 | 24 hs | 7 ds | • | • | 50-200 | 10 – 40 | 30' |

| ANTIINFECCIOSO | VIAS DE ADM. | | | PRESEN- TACION | DILUYENTES PARA RECONSTITUCIÓ | VOL. DIL (ml). | CONC. FINAL (mg/ml) | ESTABILIDAD | | VEHIC. de INF. | | VOL de INF. (ml) | CONCENT. INFUSIÓN (mg/ml) | TIEMPO de ADMINIST |
|--|--------------|-------|-------|----------------------|-------------------------------------|-------------------|---------------------------|-------------|--------|----------------|-------|---------------------|---------------------------------|--------------------------|
| | IM | IVdir | IVinf | | | | | 25 °C | 2-8 °C | SF | Dx 5% | | | |
| CEFOTAXIMA | • | | | FA 1g | AD | 3,0 | 300 | 24 hs | 10 ds | | | | | |
| | | • | | FA 1g | AD | 10,0 | 95 | 24 hs | 10 ds | | | | | 3-5' |
| | | | • | FA 1g | AD | 10,0 | 95 | 24 hs | 10 ds | • | • | 50-100 | 10 – 20 | 30' |
| CEFTAZIDIMA | • | | | FA 1g | AD | 3,0 | 280 | 18 hs | 7 ds | | | | | |
| | | • | | FA 1g | AD | 10,0 | 100 | 24 hs | 7 ds | | | | | 3-5' |
| | | | • | FA 1g | AD | 10,0 | 100 | 24 hs | 7 ds | • | • | 50 – 100 | 10 – 20 | 15-30' |
| CEFTRIAXONA | • | | | FA 1g | AD | 3,0 | 280 | 24 hs | 3d | | | | | |
| | | | • | FA 1g | AD | 10,0 | 100 | 3 ds | 10 ds | • | • | 50-100 | 10 – 20 | 15-30' |
| CLARITROMICINA | | | • | FA 500 mg | AD | 10,0 | 50 | 24 hs | 48 hs | • | • | 250 | 2 | 60' |
| CLINDAMICINA | • | | | Amp. 600 mg | | | | | | | | | | |
| | | | • | Amp. 600 mg | | | | | | • | • | 50 -100 | < 18 | 20 - 60' |
| CLORANFENICOL | | • | | FA 1g | AD - Dx 5% | 10,0 | 100 | 30 ds | --- | | | | | 1' |
| | | | • | FA 1g | AD - Dx 5% | 10,0 | 100 | 30 ds | --- | • | • | 50 -100 | 10 – 20 | 30' |
| COLISTINA | • | | | FA 100 mg | AD | 2,0 | 50 | 7 ds | 7 ds | | | | | |
| | | • | | FA 100 mg | AD | 2,0 | 50 | 7 ds | 7 ds | | | | | 3-5' |
| | | | • | FA 100 mg | AD | 2,0 | 50 | 7 ds | 7 ds | • | • | | | 30' |
| COTRIMOXAZOL <i>Trimetroprima - Sulfametoxazol</i> | | | • | Amp. 80 T / 400 S | | | | | | | • | 100 - 200 | 0,4 - 0,8 T | 60 - 90' |
| GANCICLOVIR | | | • | FA 500 mg | AD | 10 | 50 | 12 hs | | • | • | 50-100 | 10/05/09 | 60' |
| GENTAMCINA | • | | | Amp. 80 mg | | | | | | • | • | | | |
| | | | • | Amp. 80 mg | | | | | | • | • | 50 - 200 | Variable | 30 - 120' |

| ANTIINFECCIOSO | VIAS DE ADM. | | | PRESENTACION | DILUYENTES PARA RECONSTITUCIÓN | VOL. DIL (ml). | CONC. FINAL (mg/ml) | ESTABILIDAD | | VEHIC. de INF. | | VOL de INF. (ml) | CONCENT. INFUSIÓN (mg/ml) | TIEMPO de ADMINIST |
|----------------------------------|--------------|-------|-------|------------------|--------------------------------|----------------|---------------------|-------------|---------|----------------|-------|-------------------------|---------------------------|--------------------|
| | IM | IVdir | IVinf | | | | | 25 °C | 2-8 °C | SF | Dx 5% | | | |
| ESTREPTOMICINA | • | | | FA 1 g | AD | 4,2 | 200 | 24 hs | 48 hs * | | | | | |
| | • | | | | | 3,2 | 250 | 24 hs | 48 hs * | | | | | |
| | • | | | | | 1,8 | 400 | 24 hs | 48 hs * | | | | | |
| IMIPENEM | | | • | FA 500 mg | SF | 100 | 5 | 10 hs | 48 hs | • | | es el de reconstitución | 5 | 30 - 60' |
| | | | | | Dx 5% | 100 | 5 | 4 hs | 24 hs | | • | | | |
| MEROPENEM | | • | | FA 1g | AD | 20,0 | 50 | 8 hs | 48 hs | | | | | |
| | | | • | FA 1g | AD | 20,0 | 50 | 8 hs | 48 hs | • | • | 50 -200 | 5 - 20 | 15 -30' |
| PENICILINA G | • | | | FA 1M | AD - SF - Dx 5% | 4,5 | 200000 U | 24 hs | 7 ds | | | | | |
| | | | | FA 3M | AD - SF - Dx 5% | 9,0 | 300000 U | 24 hs | 7 ds | | | | | |
| | | | • | FA 1M | AD - SF | 4,5 | 200000 U | 24 hs | 7 ds | • | | 50 - 100 | Variable | 1 - 2 hs. |
| | | | | FA 3M | AD - SF | 9,0 | 300000 U | 24 hs | 7 ds | | | | | |
| PIPERACILINA - TAZOBACTAM | | | • | FA 4 g P / 0,5 T | AD - SF - Dx 5% | 20,0 | 200 de P | 24 hs | 48 hs | • | • | ≥ 50 | ≥ 80 de P | 30' |
| | | | | | | | | 24 hs | | | | | | |
| RIFAMPICINA | | | • | FA 600 mg | AD | 10,0 | 60 | 24 hs | --- | | • | 100 | 6 | ≥ 30' |
| | | | | | | | | 24 hs | | | | 500 | 1,2 | 3 hs. |
| TEICOPLANINA | • | | | FA 200/400 mg | AD | 3 | | 48 hs | 7 ds | | | | | |
| | | • | | FA 200/400 mg | AD | 3 | | 48 hs | 7 ds | | | | | 1' |
| | | | • | FA 200/400 mg | AD | 3 | | 48 hs | 7 ds | • | • | | | 30' |
| TIGECICLINA | | | • | FA 50 mg | SF - Dx5% - RL | 5,3 ml | 10 | 6h | | • | • | 100 | 0,5 | 30-60 |
| VANCOMICINA | | | • | FA 500 mg | AD | 10,0 | 50 | 14 ds | 14 ds | • | • | 100 - 200 | ≤ 5 | ≥ 1 h |